


Mechanisms of light emission


Laws of thermal radiation and Luminescence



Zsolt Mártonfalvi

1

Thermal radiation



Consequently:
All bodies emit radiation independently of the temperature of their surroundings. (Prévost, 1791)
The emitted radiation is always **electromagnetic radiation**.




Quantitative description of thermal radiation:

- Radiant emittance (M)**

$$M = \frac{\Delta P}{\Delta A} = \left[\frac{W}{m^2} \right]$$

- Absorption coefficient (α)**

$$\alpha = \frac{J_{\text{absorbed by the surface}}}{J_{\text{received by the surface}}} \quad (0 \leq \alpha \leq 1)$$



M and α strongly depend on the absolute temperature of the body!


2

Kirchhoff's law

Bodies that emit more also absorb more. The ratio between radiant emittance and absorption coefficient is constant within a narrow range of wavelength (λ):



$$\frac{M_{\lambda \text{ body1}}}{\alpha_{\lambda \text{ body1}}} = \frac{M_{\lambda \text{ body2}}}{\alpha_{\lambda \text{ body2}}} = \text{constant}$$



3


The ideal black body

A theoretical body that is used as a model in the description of thermal radiation. It absorbs all radiation that falls on it:

$$\alpha_{\text{black body}} = 1$$

thus, we can calculate any real body's radiant emittance if we know its absorption coefficient ($\alpha_{\lambda i}$):

$$M_{\lambda i} = \alpha_{\lambda i} M_{\lambda \text{ black body}}$$



Stefan's law:

$$M = \sigma T^4$$

$$\sigma = 5.67 \cdot 10^{-8} \frac{W}{m^2 K^4}$$



4

The ideal black body

The emission spectrum of the black body is **continuous**.

Wien's displacement law:
 $\lambda_{max}T = constant$
 $constant = 0.0029 \text{ m} \cdot \text{K}$

Max Planck's radiation law:
 The radiation energy of a black body is emitted in discrete integer multiples of a fundamental "package", the quantum.
 $E = hf$
 $h = 6.63 \cdot 10^{-34} \text{ J} \cdot \text{s}$

$$M_{(\lambda)} = \frac{2\pi h f^3}{c^2} \cdot \frac{1}{e^{\left(\frac{hf}{k_B T}\right)} - 1}$$

5

Light sources based on thermal radiation

Sun

Hot objects

$$\frac{T_1}{T_2} = \frac{\lambda_{max2}}{\lambda_{max1}}$$

$$\frac{M_1}{M_2} = \left(\frac{T_1}{T_2}\right)^4$$

Incandescent light bulb

6

Sun light

Spectrum of Solar Radiation (Earth)

Irradiance ($\text{W}/\text{m}^2/\text{nm}$)

Wavelength (nm)

$\lambda_{max} \approx 500 \text{ nm}$

5778K blackbody

Atmospheric absorption bands

H_2O , O_3 , H_2O , H_2O , CO_2 , H_2O

7

Medical applications

Telethermography

Measures the radiant emittance of thermal radiation of the human body.
 Human body: $\lambda_{max} \approx 10 \mu\text{m}$ (infrared)
 $\alpha \approx 0.95$ (95% of ideal black body)

Radiant emittance strongly depends on temperature!

Diagnosis of inflammations, cancer, circulatory defects that cause the change in local temperature.


Medical IR scanner

8

Medical applications

Non-contact thermometry

Typically measures the total radiant intensity between 8-14 μm (*spectral responsivity*).
 1 °C of temperature difference will increase the radiante emittance by ~1.5% !



9

Medical applications

Heat reflecting blankets (Space blanket)

Reflects about 80% of the radiation in the infrared range.

Emergency use



Surgical use



10

Luminescence

Spontaneous emission of a photon due to the relaxation of an excited electron.

Phases of luminescence:


- Absorption of external energy
- Excitation
- Emission of energy in the form of electromagnetic radiation

Types of luminescence

Type of excitation	Name	Example
Light	Photoluminescence	Fluorescent lamp
Electric	Electroluminescence	Na-lamp
Radioactive	Radioluminescence	Nal (Tl) (scintillator)
Mechanical	Triboluminescence	(Percussion, friction)
Biochemical	Bioluminescence	firefly
Thermal	Thermoluminescence	CaSO ₄ (Dy) (dosimeter)

Type of relaxation	Name	Example
S ₁ – S ₀ (fast)	Fluorescence	Fluorescein
T ₁ – S ₀ (slow)	Phosphorescence	Phosphorous

11



12

Luminescence of atoms

Luminescence emission of atoms has a **line spectrum**.

$E_{\text{photon}} = E_2 - E_1 = hf = h \frac{c}{\lambda_{em}}$

Flame test

Energy Levels of Hydrogen

Visible light transitions of the hydrogen atom

shells or "orbits" of electron

13

Luminescence of molecules

Discrete energy levels split into **vibrational levels**.

The energy of a molecule is the sum of its electronic, vibrational, and rotational transition energies.:

$$E_{\text{total}} = E_e + E_v + E_r$$

Molecular vibrations: (2 examples)

Luminescence of molecules has **band spectrum**

Spin states of excited electrons

Singlet state (S)
Sum of spin quantum numbers is $S = 0$ (+1/2, -1/2)

Triplet state (T)
Sum of spin quantum numbers is $S = 1$ (+1/2, +1/2)

Magnetic moment of spin state: $2S + 1$

14

Luminescence emission of molecules

Jablonski diagram

Kasha's rule: The excited molecule first reaches the lowest vibrational level of S_1 and photon emission occurs always from this state to any vibrational level of the ground (S_0) state.

Aleksander Jablonski

Michael Kasha

15

Luminescence emission of molecules

Jablonski diagram

Lifetimes (τ)

- Excitation: femtosecond (10^{-15} s)
- Vibrational relaxation: picosecond (10^{-12} s)
- Fluorescence emission: nanosecond (10^{-9} s)
- Phosphorescence emission: microsecond to seconds ($10^{-6} - 10$ s)

16

Luminescence emission of molecules

Quantum yield (Q_F):

$$Q_F = \frac{k_f}{k_f + k_{nr}} = \frac{\text{number of photons emitted}}{\text{number of photons absorbed}}$$


Lifetime (τ):

$$N = N_0 \cdot e^{-\frac{t}{\tau}}$$

$$\tau = \frac{1}{k_f + k_{nr}}$$

k_f : Rate of photon producing transitions


k_{nr} : Rate of non-radiative transitions



17

Stokes shift


The shift between the emission and excitation spectra



18

Comparing emission processes


	Fluorescence	Phosphorescence
Relaxation	From singlet state $S_1 \rightarrow S_0$	From triplet state $T_1 \rightarrow S_0$
Lifetime	Nanoseconds	From microseconds up to seconds
Stokes shift	Smaller	Larger (because T_1 is at lower energy than S_1)



19

Measurement of luminescence

Luminescence spectrofluorimeter



20

Applications - FRET

Förster Resonance Energy Transfer

Energy transfers from donor without emission to acceptor in dipole-dipole interactions. Requires spectral overlap between donor emission and acceptor absorption.

Intensity vs. Wavelength (λ_{ex} and λ_{em})

FRET efficiency or quantum yield (E):

$$E = \frac{1}{1 + (r/r_0)^6}$$

r_0 : Förster distance

"Molecular ruler"

21

Applications - FRAP

Fluorescence Recovery After Photobleaching

Photobleaching: The permanent loss of fluorescence due to photochemical reactions.

Top View vs. Side View (molecular detail, not to scale)

A, B, C, D

Normalized Fluorescence Intensity vs. Time (s)

Lateral diffusion of lipid molecules

22

Fluorescence microscope

23

Biomedical applications

Fluorescent transgenic animals

Fluorescence guided surgery

Surgeon's former view vs. Surgeon's new view

View of localized region in peritoneal cavity of an ovarian cancer patient as seen with the naked eye (left) or with the aid of a tumor-targeted fluorescence dye (right).

24