


Biophysics I

13. Nuclear radiations

Liliom, Károly

01. 12. 2023.

liliom.karoly@med.semmelweis-univ.hu
karoly.liliom.mta@gmail.com




Constituents of atoms

| Particle | Symbol | Resting Energy (MeV) | Relative Charge* | Mass (kg) | Relative Mass (AMU)** |
|----------|--------|----------------------|------------------|--------------------------|-------------------------|
| electron | e | 0.51100 | 1- | 9.11×10^{-31} | 5.4858×10^{-4} |
| proton | p | 938.272 | 1+ | 1.6726×10^{-27} | 1.0072765 |
| neutron | n | 939.566 | 0 | 1.6749×10^{-27} | 1.0086649 |


* electrons have an electric charge of $-1.602 \times 10^{-19} \text{ C}$

**The atomic mass unit is defined as 1/12 of the carbon (^{12}C) atom

Nucleus size





$1 \text{ \AA} = 100,000 \text{ fm}$



Nuclear stability

There are very large *repulsive electrostatic forces* between protons
should cause the nucleus to fly apart

It must be an attractive force be present within the nucleus!



Rutherford, 1911 – nuclear force: an attractive force acting on short distances within the nucleus, independent of charges, and stronger than the Coulomb forces.


The hypothesis of neutron (discovered by Chadwick in 1932)

Nuclear notation


Mass number

$$A = Z + N$$

*Chemical symbol for
the element*



*Atomic number =
Number of protons*



N = number of neutrons

nucleon = proton or neutron

Nuclear stability

$$\Delta M = [Zm_p + (A-Z)m_n] - M(A,Z)$$

The mass defect (or mass deficit) problem: the mass of a nucleus is less than the mass of its constituent nucleons. The difference can be explained by Einstein's law of mass-energy equivalence:


$$\Delta E = \Delta Mc^2$$

The energy corresponding to the mass defect is the binding energy of the nucleons.

Binding energy per nucleon

- The curve increases rapidly
- Sharp peaks for the even-even nuclei for ${}^4_2\text{He}$, ${}^{12}_6\text{C}$, and ${}^{16}_8\text{O}$
- Maximum is around $A=56$

nucleon = proton or neutron



Isotopes

Greek *isos topos* = *equal place*

Isotopes of an element have nuclei with

- the same number of protons
- different numbers of neutrons
- different mass number

isotope = equal place = equal atomic number


Mendeleev's Periodic Table of Elements

| | | Table of Common Polyatomic Ions | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|----|--|--------|-------|-----------------------|---------|--------|--------|--------|--------|--------|-------|------|----|-----|-----|-----|-----|------|--------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-------|
| | | Element categories | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | State of matter at 25 °C | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Gas | Liquid | Solid | Artificially prepared | Unknown | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 | IA | | | | | | 13 | 14 | 15 | 16 | 17 | 18 | VIIA | | | | | | VIIA | VIII A | | | | | | | | | | | |
| 1 | H | +1 -1 | | | | | B | C | N | O | F | He | | | | | | | | | | | | | | | | | | | |
| 2 | Li | +1 -2 | Be | | | | 10.811 | 12.011 | 14.007 | 15.999 | 18.998 | 4.003 | | | | | | | | | | | | | | | | | | | |
| 3 | Na | +1 -1 | Mg | | | | 13 | Al | Si | P | S | Cl | Ar | | | | | | | | | | | | | | | | | | |
| 4 | K | +1 -1 | Ca | | Sc | Ti | V | Cr | Mn | Fe | Co | Ni | Cu | Zn | Ga | Ge | As | Se | Br | Kr | | | | | | | | | | | |
| 5 | Rb | +1 -1 | Sr | | Y | Zr | Nb | Mo | Tc | Ru | Rh | Pd | Ag | Cd | In | Sn | Sb | Te | I | Xe | | | | | | | | | | | |
| 6 | Cs | +1 -1 | Ba | | Hf | Ta | W | Re | Os | Ir | Pt | Au | Hg | Tl | Pb | Bi | Po | At | Rn | | | | | | | | | | | | |
| 7 | Fr | +1 (223) | Ra | (226) | 104 | Rf | 105 | Db | 106 | Sg | 107 | Bh | 108 | Mt | 110 | Uun | 111 | Uub | 112 | Uut | 113 | Uuq | 114 | Uup | 115 | Uuh | 116 | Uus | 117 | Uuo | (294) |
| | | Selected Oxidation States | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Atomic Number | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Symbol | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Electron Configuration | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Atomic Mass | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | La Ce Pr Nd Pm Sm Eu Gd Tb Dy Ho Er Tm Yb Lu | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | Ac Th Pa U Np Pu Am Cm Bk Cf Es Fm Md No Lr | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Nuclear stability chart

- Light nuclei are most stable if $N=Z$
- Heavy nuclei are most stable when $N > Z$

As the number of protons increases, the Coulomb force increases and so more neutrons are needed to keep the nucleus stable
- No nucleus is stable when $Z > 83$



What does it mean „unstable”?

Radioactive decay



Antoine Becquerel
1903 Nobel Prize in Physics
for discovering radioactivity




Image of Becquerel's photographic plate which has been fogged by exposure to radiation from an uranium salt. The shadow of a metal Maltese Cross placed between the plate and the uranium salt is clearly visible. (1896)

Radioactive decay

- *Radioactivity* is the spontaneous release of energy in the form of particles or electromagnetic waves
- Experiments suggested that radioactivity was the result of the decay of unstable nuclei
- Three types of radiation can be emitted
 - Alpha (α) particles
 - Beta (β) particles
 - Gamma (γ) rays

Characteristics of radioactive decay

- it is a *statistical process* – individual disintegrations occur ***randomly***
- it results in a decrease over time of the initial number of unstable (radioactive) nuclei




Characteristics of radioactive decay

$$\text{Activity: } \Lambda = \left| \frac{\Delta N}{\Delta t} \right| \quad \begin{array}{l} \text{N: number of nuclei} \\ \text{to be decayed} \\ \text{t: time} \end{array}$$

Activity = number of nuclei decayed in a unit time


unit: becquerel (Bq) 1Bq = 1 decay/sec

background



kBq,

diagnostics




MBq,

*laboratory
practice*



GBq,

therapy



TBq

Radioactive decay law

Differential form:

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

solution

λ : decay constant, characteristic for isotopes (1/s)

Integral form:

$$N = N_0 e^{-\lambda t}$$

N_0 : number of radioactive nuclei at $t=0$,


N : number of remaining radioactive nuclei at a later time t

Activity depends both on the type of isotope and on the size of the population of unstable (radioactive) nuclei

Specific activity: activity in a unit mass of isotope (Bq/kg)


Graphical representation

$$N = N_0 e^{-\lambda t}$$





If $t = T_{1/2} \longrightarrow N_0 / 2 = N_0 e^{-\lambda T_{1/2}} \longrightarrow \lambda = \frac{\ln 2}{T_{1/2}} = \frac{0.693}{T_{1/2}}$

Half-lives in Medical Practice






Iodine - 131 (^{131}I) - $T_{1/2} = 8 \text{ days}$
Thyroid treatment

Technetium-99m ($^{99\text{m}}\text{Tc}$) – $T_{1/2} = 6 \text{ hours}$
Isotope diagnostics





Gold-198 (^{198}Au) - $T_{1/2} = 2.7 \text{ days}$
Tumor therapy



Types of radioactive decay



Types of radioactive decay



α decay



α particle is a nucleus of helium containing two neutrons and two protons



Heavy nuclei ($A > 150$) can disintegrate by emission of an α particle

example:




Energy spectrum of α radiation

example



Energy levels are characteristic for the nucleus

Penetration depth of α particles




| absorber | density | alpha range |
|---------------------|------------------------|--------------------|
| air (STP) | 1.2 mg/cm ³ | 3.7 cm |
| paper (20lb) | 0.89 g/cm ³ | 53 μ m |
| water (soft tissue) | 1.0 g/cm ³ | 45 μ m |


Medical application of α radiation

Diagnostics: none


Targeted alpha therapy of cancer



Seed implantation
by needle






Monoclonal
antibody




Carbon nano-tube

β decay



1. Neutron excess: β^- decay



example:



Energy spectrum of β radiation




*Energy distribution of the β^- particles
emitted during the β^- decay of ^{32}P .*

continuous spectrum

with a maximum kinetic energy for the β particle

Penetration depth of β^- particles




| <i>absorber</i> | <i>density</i> | <i>maximum beta range (2.3 MeV)</i> | <i>(1.1 MeV)</i> |
|------------------------|------------------------|--|-------------------------|
| air | 1.2 mg/cm ³ | 8.8 m | 3.8 m |
| water (soft tissue) | 1.0 g/cm ³ | 11 mm | 4.6 mm |
| aluminum | 2.7 g/cm ³ | 4.2 mm | 2.0 mm |
| lead | 11.3 g/cm ³ | 1.0 mm | 0.4 mm |

Medical application of β^- radiation

Diagnostics: none

Targeted therapy: hyperthyroidism, thyroid, prostate, and several other types of cancer




Brachytherapy:
implants into the
tumours



Endovascular
irradiation

β decay

2. Proton excess: β^+ decay




parent nucleus *daughter nucleus* *positron* *neutrino*



example:



Annihilation - particle-antiparticle pairs annihilate each other




1. Conservation of momentum: two photons with opposite direction are produced
2. Energy balance:

$$m_e c^2 + m_p c^2 = 2 h f$$

mass-energy equivalence


γ decay – Isomeric transition






Sometimes the newly formed isotopes (after α or β decay) appear in the excited state.

Excited nuclides release the excess of energy by emission of gamma rays.

half-life ranging from hours up to more than 600 years




Energy spectrum of γ radiation



Energy is characteristic for the nucleus


Penetration depth of γ radiation




Penetration depth is higher than that of α or β particles, but it is highly energy dependent.

Gamma rays can travel **hundreds of meters** in the air and can easily pass **through people (\sim dm)**.

Medical application of γ rays






therapy: γ -knife



*Bone scan using
 ^{99m}Tc labeled
phosphate compound*


Diagnostics: ideal for isotope diagnostics

Technetium-99m generator



$T_{1/2}=67$ hours

$T_{1/2}=6$ hours

| <u>Isotope</u> | <u>radiofarmacon</u> | <u>organ</u> | <u>function</u> |
|--------------------|--------------------------------|---|-------------------|
| $^{99}\text{Tc}^m$ | <i>sodium pertechnetate</i> | <i>brain</i> | <i>blood flow</i> |
| $^{99}\text{Tc}^m$ | <i>albumin-linked</i> | <i>lung</i> | <i>blood flow</i> |
| $^{99}\text{Tc}^m$ | <i>colloidal suspension</i> | <i>kidney</i> | <i>metabolism</i> |
| $^{99}\text{Tc}^m$ | <i>phosphate/pyrophosphate</i> | <i>bone</i> | <i>metabolism</i> |
| ^{123}I | <i>iodine</i> | <i>thyroid gland</i> | <i>metabolism</i> |
| ^{123}I | <i>hippurate</i> | <i>kidney</i> | <i>metabolism</i> |
| ^{133}X | <i>X gase</i> | <i>lung</i> | <i>breathing</i> |
| ^{18}F | <i>fluorodeoxyglucose</i> |  | <i>metabolism</i> |

Checklist

Composition and stability of the nucleus

Origin of nuclear force

Radioactive decay law – differential and integral form

Decay constant, half-life, mean life time

Types of nuclear radiation and their characteristics

Damjanovich, Fidy, Szöllősi: Medical Biophysics

I. 1.5

1.5.1

1.5.2

1.5.4

II.3.2

3.2.1

3.2.2

3.2.3

3.2.4