




Atomic and molecular interactions; applications

Tamás Bozó
30 October 2014

*Damjanovich, Fidy, Szöllösi: Medical Biophysics, pp. 44-51; 579-583
Lab Manual, Chapter 18.c., Resonance*

1




Atomic interactions

Noble gases: found in their atomic form in nature (*He, Ne, Ar, Kr, Xe, Rn*)

Other elements: form molecules held together by chemical bonds. (e.g. H_2, HCl, H_2O, \dots) Each atom has a (more or less) fixed position in the molecule.

3



Bond Types

Classification systems:

- Intra/intermolecular bonds
- According to the strength of bonds:
 - Strong/weak
 - Primary (covalent, metallic, ionic)
 - Secondary (dispersion, dipole-dipole, hydrogen)
- Other Bond types...



5

I. Covalent bond



- Atoms are held together by electrons of common orbitals, called molecular orbitals.
- Valence electrons interact the most.
- Single, double or triple bond depending on the number of shared electrons.
- Some electrostatic component may be present (see later), but not predominant.
- Strong interaction: $E_k > 1 \text{ eV}$
- ($1 \text{ eV} = 1,6 \cdot 10^{-19} \text{ J} = 96 \text{ kJ/mol} \sim 100 \text{ kJ/mol}$)


	R_0	E_b
H_3C-CH_3 ethane	154 pm	-331 kJ·mol ⁻¹
$H_2C=CH_2$ ethylene	139,9 pm	-590 kJ·mol ⁻¹
$HC\equiv CH$ acetylene	120,3 pm	-812 kJ·mol ⁻¹

6

I. Covalent bond


Apolar (nonpolar or homeopolar) bond

Center of (+) és (-) charges of the molecule coincides
 Binding electron(s) shared equally
 „purely covalent”
 E.g. H_2 , Cl_2 , O_2 , ...



Polar (heteropolar) bond

Centers of (+) and (-) charges do not coincide
 Electrostatic contribution appears
 Electron cloud is polarized
 Electric dipole is formed
 E.g.: HCl , HF , H_2O , ...




7

I. Covalent bond

Electric dipole moment: measure of the separation of positive and negative electric charges.

$$p = Qd$$

p: electric dipole moment
 Q: charge (positive and negative)
 d: distance of separation
 [D, debye] ($1D = 3,34 \cdot 10^{-30} \text{ Cm}$)



8

I./b Metallic bond


- Atoms are held together by common electrons (delocalized orbital)
- Valence electrons form continuous electron band
- Strong interaction: $E_k > 1 \text{ eV}$
- Multi atomic systems

Periodic Table of the Elements

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

9

I./b Metallic bond




Metallic lattice: positively charged metal ions in a crystall lattice surrounded by a cloud of delocalized electrons.



10

I./b Metallic bond



Physical properties:


- Metals are:
- Non transparent
 - Ductile (shapeable)
 - Good electric conductors
 - etc.

11

II. Bond types involving electrostatic interactions

Electronegativity

- is a chemical property that describes the tendency of an atom or a functional group to attract electrons towards itself.
- The higher the associated electronegativity number, the more an element or compound attracts electrons towards it.
- Approximately proportional to the sum of ionization energy and electron affinity.
- Calculated with different methods (*Pauling, Mulliken, Sanderson...*)



12

II./b Dipole-dipole interaction

- Constant charge distribution is present in a (given part of a) molecule
- Partially (+) and (-) segments are held together by electrostatic interactions (Coulombic forces)
- Intra/intermolecular interaction.
- Weak interaction ($E_b = 0,003-0,02$ eV)
- Energy of attracting interaction between dipoles:


$$E_{\text{attraction}} = p E$$

p : dipole momentum
 E : electric field strength generated by the surrounding partners

($E_{\text{repulsion}}$: stems from the repulsion of the participant's electron cloud)

19

II./b Dipole-dipole interaction



20


III. Van der Waals-interactions

- Sum of attractive and repulsive interactions between two apolar atoms, molecules or apolar molecular parts.
- The **attractive contribution** (also called *London-*, or *dispersion force*) is a result of temporarily created dipoles that can induce the polarization of other apolar molecule or molecular part.
- Intermolecular or intramolecular interaction.
- Important biological role: formation of organic structures.
- Weak connection ($E_b \sim 0,02$ eV)
- [according to other classifications Van der Waals interactions involve all types of weak electrostatic connections (permanent dipole-permanent dipole; permanent dipole-induced dipole, induced dipole-induced-dipole)]

21

III. Van der Waals-interactions


Equilibrium distance of a Van der Waals interaction can be considered as one definition of atomic size.



22

IV. Hydrogen bond

- Two atoms of high electronegativity are held together with a Hydrogen-bridge.
- Primarily between **F, N, O** atoms (pillar atoms).
- Intermolecular/intramolecular interaction
- Typical bond distance: 0,23 – 0,35 nm
- Well defined geometry.
- Important role in structural biology and biochemical reactions.
- Medium strong interaction (typical $E_b = 0,2$ eV)



23


V. Hydrophobic interaction

- Appears in aqueous systems (like biological environments!)
- Apolar, hydrophobic molecules tend to aggregate (form bonds with each other) in aqueous environment in order to exclude the polar water molecules and minimize their surface area exposed to water.
- Intra/intermolecular interaction
- It has mostly entropic origin (see later at Thermodynamics) through reduction of highly structured water cage around the apolar surfaces. (see Organisation of Water later)
- Important role in structural biology and biochemical reactions.
- Weak interaction.



24

V. Hidrophobic interaction



25

Scanning Probe Microscopy (SPM)

Family of instruments used for studying surface properties of various materials.
How do they work?

Etymology and function:

Microscopy: a method being able to form image of small objects.
How small? Size of resolvable objects spans from few pm-s to several μm -s.

They are not „scope“-s in the classic sense of the word: They do not „see“ the object, they „touch“ it.

26

Scanning Probe Microscopy (SPM)

Probe:

- A tiny, specifically designed component particularly sensitive to *atomic interactions*.
- The probe is brought very closed to the sample surface.
- The sensed interactions can be correlated with the distance between the probe and the sample.
- Various interactions can be observed depending on the design of the probe.
- SPM methods are named after the type of atomic interaction sensed by a certain probe.

27


Scanning Probe Microscopy (SPM)

Scanning:


- A scanner controls the precise *position* (X; Y) of the probe and its *distance* (Z) from the surface to be imaged.
- The probe can be moved with pm sensitivity in X; Y; Z directions.
- The surface of region of interest (ROI) is scanned point by point during a measurement.
- (The material that enables such precise positioning is *piezoelectric ceramic*. If voltage is applied on it, the ceramic changes its geometry.)

28

Scanning Tunneling Microscope (STM) 1981




Atoms in a silicon chip



Heinrich ROHRER and Gerd BINNING
Nobel prize: 1986


29

Scanning Probe Microscopy "Family Tree" (SPM)



30


Atomic Force Microscopy



The first one. 1986.


31

Atomic Force Microscopy Tip at the end of a cantilever




32

Atomic Force Microscopy Different types of AFM cantilevers.



33


Atomic Force Microscopy



34

Atomic Force Microscopy

- Bringing the tip very close (few nm-s) to the sample atomic interactions arise between the very last atoms of it and the atoms of the sample.




Depending on the tip-sample distance:

- Net attraction or repulsion may occur.
- Attraction at „longer” distances.
- Repulsion at very short distances.

35

Atomic Force Microscopy



- Attractive and repulsive effects cause the cantilever to bend.
- Deflection of the cantilever is detected by an optical system.
- A laser beam is pointed at the very end of the cantilever, and reflected back to be sensed by a position-sensitive photodetector (PSD).
- Thus vertical deflection of the cantilever is amplified and can be detected with sub-Å sensitivity.

36

Atomic Force Microscopy (AFM)


- The probe is a tiny, sharp tip, usually few μm -s tall, and only few nm-s in diameter at the apex.
- The tip is located at the free end of a cantilever that is usually 100-500 μm long.
- Topographic image is collected with ~ 10 pm vertical and somewhat worse horizontal resolution.
- Any surfaces (conductors, insulators and semiconductors) can be imaged.
- Works in air and in fluid environment as well.
- Usually does not require fixation or staining of the sample.
- Biological samples can be examined in their native state and physiological environment.

37

Atomic Force Microscopy

Contact mode:

- The tip is in perpetual contact with the sample surface.
- The deflection of the cantilever, i.e. the force exerted on the sample by the tip is held constant.
- A Z feedback system is utilized to maintain the deflection at a constant value (setpoint) by lifting or lowering the cantilever.
- Topography data (i.e.: height) in each X;Y point is calculated from these Z movements



Disadvantage:


- Lateral forces exerted by the scanning tip may damage softer samples.

38

Atomic Force Microscopy

Oscillating mode: (Tapping mode, Non-contact mode)

- Cantilever is oscillated close to its resonant frequency.
- The tip taps the surface gently
- The amplitude of cantilever oscillation changes with surface topography.
- A Z feedback system is utilized to maintain the amplitude at a constant value (setpoint) by lifting or lowering the oscillating cantilever.
- Topography data (i.e.: height) in each X;Y point is calculated from these Z movements




Advantage:


- Virtually eliminated lateral forces.
- Allows more gentle imaging.
- Applicable for soft samples.

39

Scanning Probe Microscopy (SPM)




Pentacene molecule imaged with AFM



Hydrogen bonds between 8-hydroxyquinoline molecules scanned with AFM

40

Atomic Force Microscopy Images




NaCl crystal surface

Human metaphase chromosomes

41

Atomic Force Microscopy Images



„The thinker“ a single actin polymer

HeLa cells on glass

42




Atomic Force Microscopy

<http://www.youtube.com/watch?v=BrsoS5e39H8>

43

Atomic Force Microscopy


Force spectroscopy: Force-distance traces registered upon push-pull cycles of the AFM tip (movement only in Z direction)
~10 pN sensitivity



44

Atomic Force Microscopy

Force spectroscopy:



45

Thank you for your attention!



Fabio Picasso: "Don Quixote"
polikarbonát felszínre rajzolva

46