

Mathematical and Physical Basis of Medical Biophysics

Chapter 5 Mechanics –Work and Energy

Judit Somkuti 16/09/2016

Energy-Work

Work and energy describes the interactions between objects, can be applied more widely than force (e.g. for thermal or chemical interaction)

Energy:

- Describes the state of an object or a system
- Ability of a system to perform work

Work: is done on an object when you transfer energy to that

Different kinds of energies

- Kinetic energy
 - Potential energy
 - Internal energy
 - Chemical energy
 - Nuclear energy
 - Electrical energy
- ... they can be converted into each other

Work

For mechanical interactions:

- Object accelerated by force
- Lifting up an object
- Extending a spring



$$W = F \cdot s$$


(if force and displacement have the same direction)

W: Work [N · m=Joule=J] (scalar)

F: force [N=Newton= $\frac{kg \cdot m}{s^2}$]

s: displacement of the object [m]

$$W = F \cdot s \cdot \cos \alpha$$



α : angle between the force and the displacement

Power

- Rate of doing work

$$P = W/t$$

P: power [J/s=watt=W]

W: Work [N ·m=Joule=J]

t: time [s]

Energy and work have the same unit: J


Unit conversions

| | Joule | Electronvolt | Calorie |
|--------|----------------------|----------------------|--------------|
| 1 J= | 1 | $6.25 \cdot 10^{18}$ | 0.239 |
| 1 eV= | $1.6 \cdot 10^{-19}$ | 1 | not relevant |
| 1 cal= | 4.19 | not relevant | 1 |

Problem V/2

A car ($m=1,2 \text{ t}$) is uniformly accelerating from rest for 12 seconds to reach a velocity of 100 km/h.

- Calculate the force necessary for this acceleration.
- Calculate the distance run by the car during acceleration.
- Calculate the work done by the accelerating force.
- Calculate the average power of the car.
- Calculate the kinetic energy of the car at the end of the acceleration.



Types of mechanical energy

- Kinetic energy
(accelerating an object)
- Gravitational potential energy
(lifting up an object)
- Elastic energy
(extending a spring)

Kinetic energy

- Related to motion

$$E_{kin} = \frac{1}{2} \cdot m \cdot v^2$$

E_{kin} : kinetic energy [J=Joule]

m: mass [kg]

v: velocity [m/s]

Work done during acceleration: $W = F \cdot s = m \frac{v}{t} \cdot \frac{v}{2} t = \frac{1}{2} m \cdot v^2$.

Potential energy

- Results from a position or configuration

Depending on the force field it can be:

- Gravitational
- Magnetic (later)
- Electric (later)

Elastic energy: configuration dependent potential energy

Gravitational potential energy

- A capacity for doing work as a result of the object position in a gravitational field


$$E_{pot} = m \cdot g \cdot h$$

E_{pot} : potential energy [J=Joule]

m: mass[kg]

g: gravitational acceleration=9.81 [m/s²]

h: height above a reference level [m]



Work done during elevation:


$$W = F \cdot s = mgh$$

Problem V/4

The left ventricle pump about 70 g of blood in a contraction into the aorta. This amount of blood reaches the aortic arch that is located approximately 15 cm above the ventricle and has a flow velocity of 30 cm/s.

Calculate:


- a) the work needed to lift the blood,
- b) the work needed to accelerate the blood.
- c) the power of the left ventricle during a contraction that lasts for 0,2 s!



Problem V/5


Someone is pulling a bucket full of water ($m=12$ kg, including the mass of 10 litres of water in it) to the top of an 8 m deep well, with uniform velocity of 50 cm/s. Calculate

- the force acting on the bucket
- the work done
- the power
- How many kcal of energy equals the work of a man who is lifting up a total of $4,8 \text{ m}^3$ water from the well during one day?




Problem V/9

A ball ($m=0,8$ kg) hits the floor from a height of 2 m and bounces back to the height of 1,2 m. Calculate the amount of energy lost due to air drag and the collision with the ground.



Elastic energy



$$E_{elastic} = \frac{1}{2} \cdot k \cdot s^2$$

$E_{elastic}$: elastic potential energy [J=Joule]

k: spring constant [N/m]

s: deformation of an elastic object [m]




Work done during the stretch:

$$W = F \cdot s = \frac{1}{2} k s \cdot s = \frac{1}{2} k s^2$$

Problem V/7

Calculate the amount of energy stored in the Achilles tendon with a spring constant of $3 \cdot 10^5$ N/m that is extended by 2mm.



Conservation of energy for mechanics

- Total amount of energy in an isolated system remains constant
- Isolated system: neither matter nor energy can pass

$$E_{kin} + E_{pot} + E_{elastic} = constant$$

$$\frac{1}{2} \cdot m \cdot v^2 + m \cdot g \cdot h + \frac{1}{2} \cdot k \cdot s^2 = constant$$



Mass-energy equivalence

- Every object of mass m , has a rest energy:

$$E = mc^2$$

E : rest energy [J=Joule]

m : mass [kg]


c : speed of light in vacuum $= 3 \cdot 10^8$ [m/s]

Mass and energy can be transferred into each other.

e.g. PET

Problem V/10

Calculate the rest energy of an electron ($m_e = 9,11 \cdot 10^{-31}$ kg)! Convert your result to eV unit !




Chapter 6

Mechanics –Pressure

Pressure

$$p = \frac{F}{A}$$



p : pressure [$\text{N}/\text{m}^2 = \text{Pa} = \text{pascal}$]

F : force applied perpendicular to the surface [N]

A : surface of an object [m^2]

Units of pressure

SI unit: pascal ($\text{Pa} = \text{N}/\text{m}^2$)

Other units:

$$1 \text{ bar} = 10^5 \text{ Pa} = 100 \text{ kPa}$$

$$1 \text{ atm} = 1.01 \cdot 10^5 \text{ Pa} = 101 \text{ kPa} = 1.01 \text{ bar} = 760 \text{ mmHg}$$

$$1 \text{ mmHg} = 1 \text{ torr} = 133 \text{ Pa} = 0.133 \text{ kPa}$$

Problem VI/2

Masticatory forces of a human can reach up to 100 N (for crocodiles it is 1000 N!). When someone bites on a bone chip in the burger or on the seed of a fruit, this force is concentrated on a surface area of 1 mm^2 . Calculate the pressure!



Problem VI/3

- Calculate the pressure that a 70 kg standing man exerts on the floor. (The total surface of the two soles is 200 cm^2)
- Calculate the pressure this man exerts on the ice surface during skating! (The total surface of the blades is 4 cm^2)



Density

- Pressure in gases and fluids depends on density

$$\rho = \frac{m}{V}$$

if it is homogenous

ρ : density [kg/m³]

m: mass [kg]

V: volume [m³]

| Density of materials | |
|------------------------------|-----------------------------|
| material | ρ (g/cm ³) |
| air (at 0°C and 101 kPa) | 0.00129 |
| water (at 4°C and 101 kPa) | 1 |
| water (at 100°C and 101 kPa) | 0.958 |
| ice | 0.92 |
| aluminium | 2.7 |
| mercury | 13.6 |
| gold | 19.3 |
| human body (averaged) | 1.04 |

$$1\text{g/cm}^3 = 1\text{kg/dm}^3 = 1000\text{ kg/m}^3$$

$$\text{ml} = \text{cm}^3 \quad \text{liter} = \text{dm}^3$$

Problem VI/5

- Calculate the mass of a gold cube with the width of 10 cm!
- Calculate the pressure exerted by this cube on a horizontal shelf, that holds it!

