


Atomic and molecular interactions

As an example: atomic force microscopy


(Textbook chapters: I/1.1, I/1.2, I/1.3, I/1.4, I/2, X/2 Related practice: Resonance)




Tamás Bozó

Nanobiotechnology and Molecular Biophysics Workgroup
Department of Biophysics and Radiation Biology

27 October 2017



Atomic structure




energy levels (shells) with
K: max. 2 e⁻
L: max. 8 e⁻
M: max. 18 e⁻

nucleus, including nucleons:
protons (p⁺)
neutrons (n⁰)


chemical properties!

Z: atomic number = number of protons (= number of electrons)
N: neutron number
A: mass number = Z+N
(Nuclear structure will be detailed in Lecture 11.)



Fundamental interactions in physics


Interaction	acts on	effective range	relative strength
Gravity	any particle	infinite (~1/r ²)	10 ⁻⁴⁰
Electromagnetic	charged particles	infinite (~1/r ²)	10 ⁻²
Strong nuclear	nucleons	10 ⁻¹⁵ m	1
Weak nuclear	any particle	10 ⁻¹⁸ m	10 ⁻¹³



II. Bond types involving electrostatic interactions

Electronegativity


- is a chemical property that describes the tendency of an atom or a functional group to attract electrons towards itself.
- The higher the associated electronegativity number, the more an element or compound attracts electrons towards it.
- Approximately proportional to the sum of ionization energy and electron affinity.
- Calculated with different methods (*Pauling, Mulliken, Sanderson...*)



13

II. Bond types involving electrostatic interactions

Electronegativity according to L. Pauling (dimensionless units)




14

II. Bond types involving electrostatic interactions

Classic bonds classified according to electronegativity:


$\Delta EN < 0.6$ (apolar covalent) $0.6 - 2.1$ (polar covalent) $2.1 <$ (ionic)



15

II. Bond types involving electrostatic interactions

Classic bonds classified according to electronegativity: an example



(This model utilizes Norman (and not Pauling) EN values.)


16

II./a Ionic bond

- Atoms are held together by Coulombic forces between (+) and (-) point charges
- „Limiting case of heteropolar bonds”
- Formed between atoms of significantly different electronegativity (e.g.: NaCl , $\Delta EN = 3 - 0.9 = 2.1$)
- Can form between two atoms, but ions are usually multi-atom systems.
- Long range interaction - attraction is inversely proportional to the distance (decreases slowly with it).
- Electrostatic interaction can be largely affected by other charged components (eg. *dissociation in water!*)
- Strong interaction ($E_b > 1 \text{ eV}$)

17

II./a Ionic bond




Ionic crystals: stoichiometric ratio of positive and negative ions are structured into a periodic crystalline structure. (e.g.: NaCl)



18

II./a Ionic bond



Ionization energy: is the energy required to remove electrons from gaseous atoms or ions.

Electron affinity: amount of energy released when an electron is added to a neutral atom or molecule to form a negative ion (measured in the gaseous state).

Lattice energy: measure of the strength of bonds in an ionic compound. Energy required to completely separate one mole of a solid ionic compound into gaseous ionic constituents.

19

II./b Dipole-dipole interaction

- Constant charge distribution is present in a (given part of a) molecule
- Partially (+) and (-) segments are held together by electrostatic interactions (Coulombic forces)
- Intra/intermolecular interaction.
- Weak interaction ($E_b = 0.003\text{-}0.02 \text{ eV}$)


- Energy of attracting interaction between dipoles:

$$E_{\text{attraction}} = p \cdot E$$

p : dipole momentum
 E : electric field strength generated by the surrounding partners

20

II./b Dipole-dipole interaction



21


III. Van der Waals-interactions

- Sum of attractive and repulsive interactions between two apolar atoms, molecules or apolar molecular parts.
- The **attractive contribution** (also called *London-*, or *dispersion force*) is a result of temporarily created dipoles that can induce the polarization of other apolar molecule or molecular part.
- Intermolecular or intramolecular interaction.
- Important biological role: formation of organic structures.
- Weak connection ($E_b \sim 0.02 \text{ eV}$)
- [according to other classifications Van der Waals interactions involve all types of weak electrostatic connections (permanent dipole-permanent dipole; permanent dipole-induced dipole, induced dipole-induced-dipole)]

22

III. Van der Waals-interactions


Equilibrium distance of a Van der Waals interaction can be considered as one definition of atomic size.



23

IV. Hydrogen bond

- Two atoms of high electronegativity are held together with a Hydrogen-bridge.
- Primarily between **F, N, O** atoms (pillar atoms).
- Intermolecular/intramolecular interaction
- Typical bond distance: 0.23 – 0.35 nm
- Well defined geometry.
- Important role in structural biology and biochemical reactions.
- Medium strong interaction (typical $E_b = 0.2 \text{ eV}$)



24


V. Hidrophobic interaction

- Appears in aqueous systems (like biological environments!)
- Apolar, hydrophobic molecules tend to aggregate (form bonds with each other) in aqueous environment in order to exclude the polar water molecules and minimize their surface area exposed to water.
- Intra/intermolecular interaction
- It has mostly entropic origin (*see later at Thermodynamics*) through reduction of highly structured water cage around the apolar surfaces. (*see Organisation of Water later*)
- Important role in structural biology and biochemical reactions.
- Weak interaction.



25

V. Hidrophobic interaction



26

Scanning Probe Microscopy (SPM)

Family of instruments used for studying surface properties of various materials.

How do they work?

Etimology and function:

Microscopy: a method being able to form image of small objects.
How small? Size of resolvable objects spans from few pm-s to several μm -s.


They are not „scope“-s in the classic sense of the word: They do not „see“ the object, they „touch“ it.

27

Scanning Probe Microscopy (SPM)

Probe:

- A tiny, specifically designed component particularly sensitive to *atomic interactions*.
- The probe is brought very closed to the sample surface.
- The sensed interactions can be correlated with the distance between the probe and the sample.
- Various interactions can be observed depending on the design of the probe.
- SPM methods are named after the type of atomic interaction sensed by a certain probe.



28


Scanning Probe Microscopy (SPM)

Scanning:


- A scanner controls the precise *position* (X; Y) of the probe and its *distance* (Z) from the surface to be imaged.
- The probe can be moved with pm sensitivity in X; Y; Z directions.
- The surface of region of interest (ROI) is scanned point by point during a measurement.
- (The material that enables such precise positioning is *piezoelectric ceramic*. If voltage is applied on it, the ceramic changes its geometry. *See details in Ultrasound lecture, 2nd semester*)

29

Scanning Tuneling Microscope (STM) 1981









Atoms in a silicon chip



Heinrich ROHRER and Gerd BINNING
Nobel prize: 1986

30



Atomic Force Microscopy (AFM)


- Topographic image is collected with ~10 pm vertical and somewhat worse horizontal resolution.
- Any surfaces (conductors, insulators and semiconductors) can be imaged.
- Works in air and in fluid environment as well.
- Usually does not require fixation or staining of the sample.
- Biological samples can be examined in their native state and physiological environment.

37

Atomic Force Microscopy

Contact mode:

- The tip is in perpetual contact with the sample surface.
- The deflection of the cantilever (i.e. the force exerted on the sample by the tip) is held constant.
- A Z feedback system is utilized to maintain the deflection at a constant value (setpoint) by lifting or lowering the cantilever.
- Topography data (i.e.: height) in each X;Y point is calculated from these Z movements



Disadvantage:


- Lateral forces exerted by the scanning tip may damage softer samples.

38

Atomic Force Microscopy

Oscillating mode: (Tapping mode, Non-contact mode)

- Cantilever is oscillated close to its resonant frequency.
- The tip taps the surface gently
- The amplitude of cantilever oscillation changes with surface topography.
- A Z feedback system is utilized to maintain the amplitude at a constant value (setpoint) by lifting or lowering the oscillating cantilever.
- Topography data (i.e.: height) in each X;Y point is calculated from these Z movements




Advantage:

- Virtually eliminated lateral forces.
- Allows more gentle imaging.
- Applicable for soft samples.


39

Scanning Probe Microscopy (SPM)



Pentacene molecule
imaged with AFM

Nature Chemistry **1**, 597 - 598 (2009)




Hydrogen bonds between 8-hydroxyquinoline molecules scanned with AFM


Science, 2013; **342** (6158), 611-614

40

Atomic Force Microscopy Images




NaCl crystal surface




Human metaphase chromosomes

41

Atomic Force Microscopy Images



„The thinker“
a single actin polymer



HeLa cells on glass

42


Atomic Force Microscopy

<http://www.youtube.com/watch?v=BrsoS5e39H8>

43

Atomic Force Microscopy


Force spectroscopy: Force-distance traces registered upon push-pull cycles of the AFM tip (movement only in Z direction)
~10 pN sensitivity



44

Atomic Force Microscopy

Force spectroscopy:



Deflection of the cantilever (Δx) is proportional to the force (F) (Hooke's law):

$$F = k \cdot \Delta x$$

k : spring constant of the cantilever

Binding forces, viscous and elastic properties can be measured perturbing the sample with the tip and registering the force response.

45

Thank you for your attention!



46