

Operation of molecular motors

Szabolcs Osváth

Semmelweis University

Properties of nanosystems

- Certain properties are quantized
- Gravity is negligible compared to the other forces
- Surface/mass ratio is very large
- Dissipation of heat is virtually instantaneous
- The role of different binding forces is large
- Molecular level recognition
- Self-organization
- The effect of inertia is negligible
- Increased role of viscosity and thermal movement

Properties of biological motors





- Made of soft material (biopolymers)
- Function is governed by non-covalent forces
- There are no smooth frictionless surfaces
- Fuel: ATP (NADH), transmembrane gradient
- Work in solutions – high viscosity
- Their power is: 100-1000 ATP/s, i.e. $10^{-16} - 10^{-17}$ W
- The power of Brown motion is: 10^{-8} W
- The working principle uses Brown motion

Can the role of the Brownian motion be eliminated?

The importance of the Brownian motion is the consequence of the size and not of the experimental conditions!

The effect of the Brownian motion can not be eliminated.
It does not help if:

- Pressure is decreased
– (less frequent collisions, but smaller viscosity)
- Temperature is decreased
– (the magnitude of the fluctuations is proportional to the square root of the absolute temperature)




Function of the Brownian ratchet

- Random mixing
- Energy source
- Asymmetric energy surface along the reaction coordinate


Pulsating ratchet

The potential should be turned off for a short time only, otherwise the transport happens with equal probabilities in both directions.







Kay et al. (2007) *Angev. Chem.* 46: 72

Flashing ratchet



Temperature ratchet





- ### The working principles of biological motors
- Break the isotropy of the transport resulting from fluctuations
 - Kinetic coupling between substrate and motor
 - Substrate movement is always localized
 - Manipulating localization is important element of the transport
 - Localization is the result of non-covalent interactions

Biological application of Nonequilibrium thermodynamics

Evans-Searles FT (revision)

Evans-Searles FT for the entropy production

$$\frac{P(\Omega = A)}{P(\Omega = -A)} = e^A$$

here Ω denotes entropy production

Evans and Searles (2002) Advances in Physics, 51: 1529

Crooks FT (revision)

For a driven system in contact with thermostat:

$$\frac{P_F(A \rightarrow B)}{P_R(A \leftarrow B)} = e^{\frac{W-\Delta G}{k_B T}}$$

W is the work done when the system is driven from the state described by the value A of the control parameter into B

ΔG is the free enthalpy difference between states A and B

G. E. Crooks, J. Stat. Phys. (1998) 90: 1481

Jarzynski equality (revision)





Connects the work measured during non-equilibrium measurements to the free enthalpy difference between the initial and final states.

$$\left\langle e^{\frac{-W}{k_B T}} \right\rangle = e^{\frac{-\Delta G}{k_B T}}$$

W is the work done when the system is driven from the state described by the value A of the control parameter into B .

The transition between the initial and final states need not happen through equilibrium states.

C. Jarzynski, Phys. Rev. Lett. (1997) 78: 2690




F₁-ATP-ase flashing ratchet mechanism

Rotation mechanism of the free-rotating and the loaded F₁-ATP-ase motor.


relaxed state:
continuous line

nucleotide binding state:
dashed line




Sancho and Perez-Carrasco (2010) Fluct. Noise. Lett. 11: 1240003

Power release of the F₁-ATP-ase




Sancho and Perez-Carrasco (2010) Fluct. Noise. Lett. 11: 1240003

Efficiency of the F₁-ATP-ase



Sancho and Perez-Carrasco (2010) Fluct. Noise. Lett. 11: 1240003

Structure of bacteriorhodopsin



Oesterhelt et al. (2000) Science 288: 143

