


Modelling of ligand-protein binding

I. Computation of thermodynamic quantities

Outline

- Molecular dynamics
 - Theoretical background
- Force fields
 - Energy terms
 - Parameters
- Sampling in molecular dynamics
- Free energy difference and alchemical transformations
- Examples



Molecular dynamics

Introduction

- Molecular dynamics – link between microscopic and macroscopic quantities
 - structure
 - dynamics
 - thermodynamics



FnIII_10_preSMD.mpg

History

- Alder, B. J. and Wainwright, T. E.
J. Chem. Phys. **27**, 1208 (1957)

- McCammon, J. A., Gelin, B. R.,
and Karplus, M.
Nature (Lond.) **267**, 585 (1977)

The dynamics of a folded globular protein (bovine pancreatic trypsin inhibitor) have been studied by solving the equations of motion for the atoms with an empirical potential energy function. The results provide the magnitude, correlations and decay of fluctuations about the average structure. These suggest that the protein interior is fluid-like in that the local atom motions have a diffusional character.

relatively free diffusion, while in the low $g(r)$ states diffusion is much restricted.

The conjecture¹ that some high-order virial coefficients might be negative is not necessarily supported by the present results, since only to the left of the apparent transition do the latter give lower pressures than the five-term virial expression.

Some further investigation for both 32 molecule and larger systems will be made on the present calculators, but a satisfactory determination of the detailed behavior in the apparent transition region will require higher speed equipment. The possibility that a similar phenomenon for hard spheres in two dimensions may have been missed in the original Monte Carlo calculations¹ will also be investigated.

* Work performed under the auspices of the U. S. Atomic Energy Commission.

¹ M. N. Rosenbluth and A. W. Rosenbluth, *J. Chem. Phys.* **22**, 881 (1954).

² Metropolis, Rosenbluth, Rosenbluth, Teller, and Teller, *J. Chem. Phys.* **21**, 1087 (1953).

³ B. J. Alder and T. Wainwright, *J. Chem. Phys.* **27**, 1208 (1957).

⁴ W. W. Wood and F. R. Parker, *J. Chem. Phys.* **27**, 720 (1957).

This paper discusses the Monte Carlo method in some detail, as well as giving computational results for Lennard-Jones molecules.

⁵ Kirkwood, Maun, and Alder, *J. Chem. Phys.* **18**, 1040 (1950).

Phase Transition for a Hard Sphere System

B. J. ALDER AND T. E. WAINWRIGHT

University of California Radiation Laboratory, Livermore, California

(Received August 12, 1957)

A CALCULATION of molecular dynamic motion has been designed principally to study the relaxations accompanying various nonequilibrium phenomena. The method consists of solving exactly (to the number of significant figures carried) the simultaneous classical equations of motion of several hundred particles by means of fast electronic computers. Some of the details as they relate to hard spheres and to particles

A 32-particle system in a cube and initially in a face-centered cubic lattice proceeded at about 300 collisions an hour on the UNIVAC. For comparison a 96-particle system in a rectangular box and initially in a hexagonal arrangement has been calculated, however only at high densities so far. No differences in the pressures can be detected. It became apparent that some long runs were necessary at intermediate densities, accordingly the IBM-704 was utilized where, for 32 particles, an hour is required for 7000 collisions. Larger systems of 108, 256, and 500 particles can also conveniently be handled; in an hour 2000, 1000, and 500 collisions, respectively, can be calculated. The results for 256 and 500 particles are not now presented due to inadequate statistics.

The equation of state shown in Fig. 1 of the accompanying paper¹ for 32 and 108 particles is for the intermediate region of density, where disagreement was found with the previous Monte Carlo results. The volume, v , is given relative to the volume of close packing, v_0 . Plotted also are the more extended Monte Carlo results; the agreement between these three systems is within the present accuracy of the pressure determination. This agreement provides an interesting confirmation of the postulates of statistical mechanics for this system.

Figure 1 of the accompanying paper shows two separate and overlapping branches. In the overlapping region the system can, at a given density, exist in two states with considerably different pressures. As the calculation proceeds the pressure is seen to jump suddenly from one level to the other. A study of the positions of the particles reveals that as long as the system stays on the lower branch of the curve the particles are all confined to the narrow region in space determined by their neighbors, while on the upper branch of the curve the particles have acquired enough freedom to exchange with the surrounding particles. Since the spheres are originally in ordered positions, the system starts out on the lower branch; the first jump to the upper branch can require very many collisions. The trend, as expected, is that at higher densities more collisions are necessary for

nature

SEARCH JOURNAL

Go

Journal Home
Current Issue
AOP
Archive

THIS ARTICLE

Download PDF
References

Export citation
Export references

article

Nature **267**, 585 - 590 (16 June 1977); doi:10.1038/267585a0

Dynamics of folded proteins

J. ANDREW MCCAMMON, BRUCE R. GELIN & MARTIN KARPLUS

, Cambridge, Massachusetts 02138

...bular protein (bovine pancreatic trypsin inhibitor) have been studied by solving the equations of empirical potential energy function. The results provide the magnitude, correlations and decay of ge structure. These suggest that the protein interior is fluid-like in that the local atom motions have

my account | e-alerts | subscribe | register

Tuesday 24 November 2015

Basic terms

- Thermodynamic/Macroscopic state
 - The system is characterized by few macroscopic parameters; e.g.: T , P , N
- Microscopic state
 - The system is characterized by the positions and momenta of atoms (phase space).
- Ensemble
 - Microscopic states corresponding to a macroscopic state
- Molecular dynamics simulations
 - Generation of microscopic states of an ensemble as a function of time

Thermodynamic ensembles

- Microcanonical – NVE (isolated system)
- Canonical – NVT (thermal equilibrium)
- Isotherm-izobar – NPT
- Grand canonical – μ VT (equilibrium with a reservoir of particles)

Ergodic hypothesis

- Experiment: measure the average for $\sim 10^{23}$ molecules at a certain time
- MD: calculate the time average for one (few) molecule(s)
- Measurable quantities: ensemble average $\langle A \rangle_{ensemble}$
 - e.g. (non-covalent) binding of two molecules in solution
- Molecular dynamics: time average $\langle A \rangle_{time}$

$$\langle A \rangle_{time} = \langle A \rangle_{ensemble}$$

- „long enough” MD – appropriate sampling

Force field

- Quantum mechanics
 - basic entities: nuclei and electrons
 - accurate
 - time intensive computations
- Molecular mechanics
 - basic entities: atoms
 - „classical”
 - simple, fast computations
 - includes parameters
 - Can be applied within the validity of the parameter space
 - Chemical reactions are typically outside the validity

Molecular mechanics

Atoms are pointlike objects with mass and interactions

$$E = E_{\text{str}} + E_{\text{bend}} + E_{\text{tors}} + E_{\text{vdw}} + E_{\text{el}} + E_{\text{cross}}$$


intramolecular

intermolecular

Bond stretching energy

$$E_{str} = k(r - r_0)^2$$


$$F_{str} = -2k(r - r_0)$$



good approximation in the vicinity of r_0
 k and r_0 are atom dependent parameters

Bond stretching energy - parameters

$$E_{str} = k(r - r_0)^2$$




general class	atom type ₁	description
<i>hydrogen types</i>		
	H	amide or imino hydrogen
	HC	explicit hydrogen attached to carbon
	HO	hydrogen on hydroxyl oxygen
	HS	hydrogen attached to sulfur
	HW	hydrogen in water
	H2	amino hydrogen in NH ₂
	H3	hydrogen of lysine or arginine (positively charged)
<i>all-atom carbon types₂</i>		
	C	sp ² carbonyl carbon and aromatic carbon with hydroxyl substituent in tyrosine
	CA	sp ² aromatic carbon in 6-membered ring with 1 substituent
	CB	sp ² aromatic carbon at junction between 5- and 6-membered rings
	CC	sp ² aromatic carbon in 5-membered ring with 1 substituent and next to a nitrogen
	CK	sp ² aromatic carbon in 5-membered ring between 2 nitrogens and bonded to 1 hydrogen (in purine)
	CM	sp ² same as CJ but one substituent
	CN	sp ² aromatic junction carbon in between 5- and 6-membered rings
	CQ	sp ² carbon in 6-membered ring of purine between 2 NC nitrogens and bonded to 1 hydrogen
	CR	sp ² aromatic carbon in 5-membered ring between 2 nitrogens and bonded to 1 H (in his)
	CT	sp ³ carbon with 4 explicit substituents
	CV	sp ² aromatic carbon in 5-membered ring bonded to 1 N and bonded to an explicit hydrogen
	CW	sp ² aromatic carbon in 5-membered ring bonded to 1 N-H and bonded to an explicit hydrogen
	C*	sp ² aromatic carbon in 5-membered ring with 1 substituent

Bond Stretching Potential Parameters			
Bond	used for	K _r / kcal mol ⁻¹ Å ⁻²	R _{eq} / Å
CT-CT	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	310.0	1.526
CT-H1	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	340.0	1.090
CT-HC	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	340.0	1.090
CT-NA	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	337.0	1.475
CR-H5	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	367.0	1.080
CR-NA	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	477.0	1.343
CW-H4	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	367.0	1.080
CW-NA	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	427.0	1.381
CW-CW	<i>BMT</i> ⁺ , <i>EMI</i> ⁺	549.0	1.350
AL-Cl	<i>TCA</i> ⁻	116.1	2.170
P-F	<i>PF</i> ₆ ⁻	260.3	1.646
NN-ON	<i>NO</i> ₃ ⁻	300.0	1.260




Bending energy

$$E_{bend} = k(\theta - \theta_0)^2$$



k and θ_0 are atom dependent parameters

Torsional/dihedral angle energy





$$E_{tors} = \frac{V_n}{2} [1 + \cos(n\phi - \phi_0)]$$


V_n – barrier height
 n - periodicity

Dihedral angle (atom types)				V_n , kcal/mol	n	ϕ_0	comments
C	CT1	NH1	C	0.2000	1	180.00	! backbone phi
NH1	C	CT1	NH1	0.6000	1	0.00	! backbone ksi
CT1	C	NH1	CT1	1.6000	1	0.00	! backbone omega
CA	CA	CA	CA	3.1000	2	180.00	! Phe side chain
H	OH1	CT2	CT1	0.4200	3	0.00	! Ser side chain

~15000 parameters



van der Waals energy



<i>short range:</i>	repulsive;	$\exp(-r)$ or r^{-12}	Pauli repulsion
<i>middle range:</i>	attractive;	r^{-6}	dispersion
<i>long range:</i>	disappears		

$$E_{vdw} = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} \right] - \left[\left(\frac{\sigma}{r} \right)^6 \right]$$

$$\sigma_{ij} = \frac{1}{2}(\sigma_i + \sigma_j)$$


$$\epsilon_{ij} = \sqrt{\epsilon_i \epsilon_j}$$

$$c * \exp\left(\frac{-r}{\sigma}\right)$$

Electrostatic energy

$$V_{el} = \frac{q_i q_j}{\epsilon r_{ij}}$$

Coulomb




MM parameters

- Derivation
 - Quantum mechanical calculations
 - Experimental data
 - Extension based on analogy
- Validation by comparing computed and experimental data
 - Macromolecular structure
 - NMR data
 - Structure and energy of van der Waals complexes
- Error compensation; mutual interdependence of parameters

Quality of MM force field

- Protein structure
- DNA, RNA structure
- Conformation of organic molecules
- Ligand-protein interactions
- Structure and interactions of lipids and membranes

– ...



RMSDs of backbone atoms
from the native structure
as a function of MD
simulation time

Selected MM force fields

- Charmm (Chemistry at HARvard Macromolecular Mechanics)
- AMBER (Assisted Model Building with Energy Refinement)
- OPLS (Optimized Potentials for Liquid Simulations)
- GROMOS (GROningen Molecular Simulation)
- MMFF (Merck Molecular Force Field)

MD algorithm

Newtonian mechanics

$$r^N(r_1, r_2 \dots r_N) \quad p^N(p_1, p_2 \dots p_N)$$

$$U(\underline{r}) \quad K(\underline{p}) = \sum_i \frac{|p_i|^2}{2m_i}$$

$$H = K + U \quad \dot{r}_i = \frac{p_i}{m_i} \quad \dot{p}_i = f_i$$

Verlet algorithm

Calculation of p_i and r_i at δt time steps

$$p_i(t), r_i(t) \rightarrow p_i(t + \delta t), r_i(t + \delta t) \rightarrow f_i(t)$$

$$p_i\left(t + \frac{1}{2}\delta t\right) = p_i(t) + \frac{1}{2}\delta t \cdot f_i(t)$$


$$r_i(t + \delta t) = r_i(t) + \frac{\delta t \cdot p_i\left(t + \frac{1}{2}\delta t\right)}{m_i} \rightarrow f_i(t + \delta t)$$

$$p_i(t + \delta t) = p_i\left(t + \frac{1}{2}\delta t\right) + \frac{1}{2}\delta t \cdot f_i(t + \delta t)$$

Typical δt for simulation of biochemical systems: 1-4 fs

MD - sampling

- Microstates appear according to Boltzmann distribution
 - $\exp\left(-\frac{E}{kT}\right)$
- Simulation time is limited by computational capacity
 - Time scale for proteins: $\sim\mu\text{s}$
- Rare events with high energy barrier cannot be straightforwardly simulated



Free energy - Sampling

$$F = -kT \ln \left[h^{-3} \iint \exp \left(-\frac{E(r, p)}{kT} \right) dp dr \right] \quad (1) \text{ Formula for free energy}$$

Free energy calculation with MD sampling is problematic

phase space incomplete in (1)

positive integral

\ln function increases monotonically

negative contribution missing


F overestimated

MD - Sampling

Time scale (s)	Amplitude (Å)	Description	# MD steps (step ~ fs)
10^{-15} - 10^{-12}	0.001-0.1	Bond stretching, bond angle deformation	1 - 1000
10^{-12} - 10^{-9}	0.1-10	Protein sidechain, loop and collective motions	10^3 - 10^6
10^{-9} - 10^{-6}	1-100	Folding of small proteins	10^6 - 10^9
10^{-6} - 10^{-1}	10-100	Protein folding, Ligand-protein binding	10^9 - 10^{14}


Free energy difference

- Sampling issue hampers the calculation of F and $\Delta F = F_{\text{Bound}} - F_{\text{Free}}$
- Special techniques for calculating $\Delta F = F_B - F_A$ (A similar to B) for similar systems
 - Free energy perturbation, Thermodynamic integration
- Thermodynamic cycle: binding free energy difference ($\Delta\Delta F$) of two similar ligands is obtained from the free energy differences of similar systems




- $\Delta\Delta F = \Delta F_{\text{bind}}^A - \Delta F_{\text{bind}}^B = \Delta F_{\text{transform}}^{\text{complex A} \rightarrow \text{B}} - \Delta F_{\text{transform}}^{\text{ligand A} \rightarrow \text{B}}$
- „alchemical” transformations: $\Delta F_{\text{transform}}^{\text{complex A} \rightarrow \text{B}}$ and $\Delta F_{\text{transform}}^{\text{ligand A} \rightarrow \text{B}}$
 - 2 transformations to obtain $\Delta\Delta F$

Alchemical transformation- coupling parameter



A

$H(r,p,\lambda=0)$



B


$H(r,p,\lambda=1)$

$$H_\lambda = H(r,p,\lambda) = (1-\lambda) H_A + \lambda H_B$$

λ – coupling parameter

H_λ may be other function of λ

H_A, H_B may depend on λ




<http://www.ks.uiuc.edu/Research/namd/2.6/ug/node36.html>

- Large perturbation – important change in the environment
- Large perturbation is computationally impractical

Selected applications


process of ligand binding



PP1 molecule finds the binding site of Src kinase in a 15 μs simulation

Steered MD


- Constant velocity pulling



$$U = \frac{1}{2} k [vt - (\vec{r} - \vec{r}_0) \cdot \vec{n}]^2$$


- Constant force pulling

Titin structure and function




Unfolding and force

titin I91 (formerly I27) domain




Constant velocity pulling

AFM – linear chain of I91 domains




Rief et al. (1997 Science 276 1109)

SMD – single I91 domain




Unfolding and structure




Lu, H., and Schulten, K. (2000).. Biophys. J. 79, 51–65

Titin Fn domain unfolding mechanism



A78
 $v=1\text{\AA}/\text{ns}$



Example – binding to lysozyme




Figure 6. Comparison of the HREX-SRTI relative binding free energy predictions to experiment

J. Chem. Theor. Comput. 2011, 7 3001

Example – neuraminidase inhibitors 1




Figure 2. Representative structures for zanamivir (A, B and C) and oseltamivir (D, E and F) bound to WT and mutant NAs from the SRSM/HREX simulations. Salt-bridges and hydrogen bonds are depicted as magenta and orange dashed lines, respectively. Positively charged, negatively charged, and uncharged polar groups are noted as blue, red, and purple circles, respectively, and residues of interest are labeled. Mutated residues are underlined.
doi:10.1371/journal.pcbi.1002665.g002

Example – neuraminidase inhibitors 2

Table 1. Comparison of experimental $\Delta\Delta G$ in oseltamivir and zanamivir for three NA mutations with estimates obtained using different computational approaches.

Method	H274Y		N294S		Y252H		RMSE
	$\Delta\Delta G$, kcal/mol		$\Delta\Delta G$, kcal/mol		$\Delta\Delta G$, kcal/mol		(RMSD), kcal/mol
	zanamivir	oseltamivir	zanamivir	oseltamivir	zanamivir	oseltamivir	
Experimental ^d	0.4 (0.1)	3.3 (0.2)*	1.2 (0.1)*	2.6 (0.2)*	0.1 (0.2)	−1.4 (0.1)	N/A (0.2)
SRMM	−5.8 (7.4)	0.7 (7.0)	8.2 (7.7)	5.8 (6.2)	−0.1 (8.7)	−0.9 (7.4)	4.2 (7.4)
SRSM	1.7 (2.9)	1.2 (3.0)	0.6 (2.0)	1.7 (1.9)	1.5 (1.7)	0.5 (1.5)	1.5 (2.2)
SRSM/HREX	1.3 (0.8)	4.1 (2.4)	2.3 (0.4)	2.2 (0.9)	0.6 (0.8)	0.7 (1.4)	1.1 (1.1)
MM-GBSA	6.2 (8.1)	0.9 (3.8)	5.7 (6.1)	−5.9 (3.6)	2.1 (2.9)	−1.9 (3.0)	4.8 (4.6)
MM-PBSA	8.4 (10.1)	3.0 (3.9)	5.8 (4.5)	−4.7 (3.2)	2.8 (3.1)	0.2 (2.6)	5.0 (4.6)
Rosetta	−0.4 (0.5)	0.8 (0.4)	−0.4 (0.3)	0.3 (0.2)	−0.1 (0.4)	0.0 (0.0)	1.7 (0.3)

^aValues were derived from the data reported by Collins et al [10].

Standard deviations are shown in parentheses. Root mean squared error (RMSE) and the RMS Standard Deviation (RMSD) are provided.

* indicates experimentally determined drug resistant mutation. 'N/A' stands for *not applicable*.

doi:10.1371/journal.pcbi.1002665.t001

PLOS Comp Biol 2012, 8 e1002665

Example – FKBP12-ligand

Standard binding free energy
Double decoupling
FEP
Free energy components




Figure 4. FKBP12 bound with ligand #8 studied previously.^{40,42} The gray parts are treated as a mean-field approximation with generalized solvent boundary potential.⁸⁶ See ref 42 for computational details.

$\Delta\Delta G_{\text{rep}}$	$\Delta\Delta G_{\text{dis}}$	$\Delta\Delta G_{\text{elec}}$	$\Delta\Delta G_{\text{c}}$	$\Delta\Delta G_{\text{t}}^{\circ}$	$\Delta\Delta G_{\text{r}}$	$\Delta G_{\text{bind}}^{\circ}$	exptl
-1.1	-21.1	-3.7	6.9	3.4	5.4	-10.2	-10.9

MD scope and limitation

- Scope
 - Structural study; structure refinement
 - Dynamics
 - conformations, ligand-protein binding, steered processes,...
 - Thermodynamics
 - Free energy changes
 - solvation, ligand-protein binding,...
- Limitations
 - Sampling
 - Accuracy of force field
 - Chemical reactions cannot be routinely studied